

Repetitions in beta-integers

L. Balková, K. Klouda, E. Pelantová *

*Doppler Institute and Department of Mathematics FNSPE,
Czech Technical University in Prague,
Trojanova 13, 120 00 Praha 2, Czech Republic*

Abstract

Classical crystals are solid materials containing arbitrarily long periodic repetitions of a single motif. In this Letter, we study the maximal possible repetition of the same motif occurring in β -integers – one dimensional models of quasicrystals. We are interested in β -integers realizing only a finite number of distinct distances between neighboring elements. In such a case, the problem may be reformulated in terms of combinatorics on words as a study of the index of infinite words coding β -integers. We will solve a particular case for β being a quadratic non-simple Parry number.

1 Introduction

This Letter takes up the study of β -integers initiated by the investigation of their asymptotic properties in [1]. Similarly as in the previous Letter, we restrict our consideration to β -integers realizing only a finite number of distinct distances between neighbors; β is then called a Parry number. For Parry numbers, the set of β -integers forms a discrete aperiodic Delone set with a self-similarity factor β and of finite local complexity. It follows herefrom that β -integers are suitable for modeling materials with aperiodic long range order, the so-called quasicrystals [2]. Classical crystals are solid materials containing arbitrarily long periodic repetitions of a single motif. Quasicrystals do not share this property.

In this Letter, we are interested in the maximal possible repetition of one motif occurring in β -integers. It turns out to be suitable to reformulate and study this problem in terms of combinatorics on words.

For Parry numbers, coding distinct distances between neighboring nonnegative β -integers with distinct letters, one obtains a right-sided infinite word u_β over a finite alphabet. The reformulation of our task in the language of combinatorics on words has the following reading: For a given factor w of the infinite word $u = u_\beta$, find the longest prefix v of the infinite periodic word $w^\omega = wwwwww \dots$ such that v occurs as a factor in $u = u_\beta$. The ratio of the lengths of v and w is called the index of the factor w in $u = u_\beta$ and is denoted by $\text{ind}(w)$. Let us note that $\text{ind}(w)$ is not necessarily an integer. Denote by k the lower integer part $\lfloor \text{ind}(w) \rfloor$ of the index of w , then the word w^k , i.e., the concatenation of k words w , is usually called the maximal integer power of w .

The index of any infinite word u can be naturally defined as

$$\text{ind}(u) = \sup\{\text{ind}(w) \mid w \text{ factor of } u\}.$$

Explicit values of the index are known only for few classes of infinite words. The index of Sturmian words has been studied in many papers ([3], [4], [5], [6]), the complete solution to the problem was provided independently by Carpi and de Luca [7] and by Damanik and Lenz [8]. Recently, the index of infinite words has reinforced its importance: Damanik in [9] considers discrete one-dimensional Schrödinger operators with aperiodic potentials generated by primitive morphisms

*e-mail: l.balkova@centrum.cz, karel@kloudak.eu, edita.pelantova@fjfi.cvut.cz

and he establishes purely singular continuous spectrum with probability one provided that the potentials (infinite words) have the index greater than three. Let us stress that infinite words u_β associated with Parry numbers belong to the class of infinite words generated by primitive morphisms, too.

Here we study the index of infinite words u_β for quadratic non-simple Parry numbers β . These words are determined by integer parameters p, q , where $p > q \geq 1$. We provide an explicit formula for $\text{ind}(u_\beta)$. In the particular case of $p = q + 1$, the infinite word u_β is Sturmian and our result may be deduced also from the well-known formula for the index of Sturmian words. We have chosen the word u_β associated with quadratic non-simple Parry numbers β for our study of the index of non-Sturmian words since for such infinite words, we dispose of detailed knowledge on arithmetical properties of β -integers and combinatorial properties of the associated infinite words u_β ([10], [11]).

The Letter is organized in the following way. In Section 2, we introduce necessary notions from combinatorics on words and we cite a relevant result on the index of Sturmian words. In Section 3, we provide the background on infinite words u_β coding β -integers for β being a non-simple quadratic Parry number. In Section 4, we determine the maximal integer power occurring in u_β (Theorem 4.6). Section 5 is devoted to the index of u_β (Theorem 5.3) and to the comparison of our result with the formula for the index of Sturmian words.

2 Preliminaries

An *alphabet* $\mathcal{A}$ is a finite set of symbols, called *letters*. Throughout this paper, the binary alphabet $\mathcal{A} = \{0, 1\}$ is used. The string $w = w_1 w_2 \dots w_k$, where $w_i \in \mathcal{A}$ for each $i = 1, 2, \dots, k$, is called a *word of length k* on $\mathcal{A}$. The length of w is then k and it is denoted by $|w| = k$. The set of all finite words together with the operation of concatenation forms a monoid; its neutral element is the empty word ε . We denote this monoid $\mathcal{A}^*$. An infinite sequence $u = u_0 u_1 u_2 \dots$ of symbols from the alphabet $\mathcal{A}$ is called an *infinite word*. A finite word w is said to be a *factor* of the (finite or infinite) word v if there exists a finite word v' and a finite or infinite word v'' such that $v = v' w v''$. If v' is the empty word, then w is called a *prefix* of v , if v'' is the empty word, then w is a *suffix* of v . If $v = v' w$, then $v w^{-1} = v'$, i.e., the word $v w^{-1}$ is obtained from v by erasing its suffix w . The set of all factors of an infinite word u is said to be the *language* of u and is denoted $\mathcal{L}(u)$. An infinite word u is called *recurrent* if every of its factors occurs infinitely many times in u and u is called *uniformly recurrent* if for every of its factors w , the set of all factors in $\mathcal{L}(u)$ that do not contain w as their factor is finite. In other words, every sufficiently long element of $\mathcal{L}(u)$ contains w as its factor.

The number of factors of the infinite word u gives us insight into its variability. The function $\mathcal{C}_u : \mathbb{N} \mapsto \mathbb{N}$ that to every n associates the number of distinct factors of length n occurring in u is called the *factor complexity* of the infinite word u . An infinite periodic word $u = w w w \dots$, where w is a finite word, is usually denoted w^ω . Its factor complexity $\mathcal{C}_u$ is bounded; it is readily seen that $\mathcal{C}_u(n) \leq |w|$. Similarly, the factor complexity of an eventually periodic word $u = w' w^\omega$, where w', w are finite words, is bounded. The necessary and sufficient condition for an infinite word to be aperiodic is validity of the equation $\mathcal{C}_u(n) \geq n + 1$ for all $n \in \mathbb{N}$. Infinite aperiodic words satisfying $\mathcal{C}_u(n) = n + 1$ for all $n \in \mathbb{N}$ are called *Sturmian words*; these are thus infinite aperiodic words of the lowest possible factor complexity. Sturmian words are the best known aperiodic words; a survey on their properties may be consulted in [12]. In particular, any Sturmian word is uniformly recurrent.

For determination of the factor complexity of an infinite word u , an essential role is played by special factors. We recall that a factor w of an infinite word u over a binary alphabet $\{0, 1\}$ is called *left special* if $0w$ and $1w$ are both factors of u , w is called *right special* if $w0$ and $w1$ are both factors of u , and w is said to be *bispecial* if w is both left special and right special.

The crucial notion of our study is the index of a factor w in a given infinite word u . Let us define first the integer power of w . For any $k \in \mathbb{N}$, the k -th *power* of w is the concatenation of k words w , usually denoted w^k . Analogously for $r \in \mathbb{Q}, r \geq 1$, we call a word v the r -th *power* of the

word w if there exists a proper prefix w' of w such that

$$v = \underbrace{w \dots w}_{[r]-\text{times}} w' \quad \text{and} \quad r = [r] + \frac{|w'|}{|w|}.$$

The r -th power of w is denoted w^r .

Our aim is to find, for a given factor w of an infinite word u , the highest power of w occurring in u . We will be interested exclusively in aperiodic uniformly recurrent words. Such words contain for every factor w only r -th powers of w with bounded r , see [13]. Therefore, for aperiodic uniformly recurrent words u , it makes sense to define the index of w in u as

$$\text{ind}(w) = \max\{r \in \mathbb{Q} \mid w^r \in \mathcal{L}(u)\}.$$

The word $w^{\text{ind}(w)}$ is called the *maximal power* of w in u , the word $w^{\lfloor \text{ind}(w) \rfloor}$ the *maximal integer power* of w in u . The index of the infinite word u is defined as

$$\text{ind}(u) = \sup\{\text{ind}(w) \mid w \in \mathcal{L}(u)\}.$$

Let us remark that an aperiodic uniformly recurrent word u can have an infinite index; even among Sturmian words, one can find words with an infinite index. The language of every Sturmian word is characterized by an irrational parameter $\alpha \in (0, 1)$, called *slope*. If α is the slope of a Sturmian word u , then the word obtained by exchanging letters in u has the slope $1 - \alpha$ and has evidently the same index as u . Consequently, we may assume without loss of generality that $\alpha > \frac{1}{2}$. In order to determine the index of a Sturmian word, we need to express α in the form of its continued fraction. Since $\frac{1}{2} < \alpha < 1$, the continued fraction of α equals $[0; 1, a_2, a_3, \dots]$. The results of [7] and [8] say that the index of a Sturmian word u with slope α is equal to

$$\text{ind}(u) = 2 + \sup\left\{a_{n+2} + \frac{q_n - 2}{q_{n+1}} \mid n \in \mathbb{N}\right\}, \quad (1)$$

where q_n is the denominator of the n -th convergent of α .

Now, let us describe a large class of uniformly recurrent words: fixed points of primitive morphisms. This class includes infinite words u_β associated with Parry numbers β . The map $\varphi : \mathcal{A}^* \mapsto \mathcal{A}^*$ is called a *morphism* if $\varphi(wv) = \varphi(w)\varphi(v)$ for every $w, v \in \mathcal{A}^*$. One may associate with φ the *morphism matrix* M_φ satisfying

$$(M_\varphi)_{ab} = \text{number of letters } b \text{ occurring in } \varphi(a),$$

for any pair of letters $a, b \in \mathcal{A}$.

Knowing for any $a \in \mathcal{A}$ the number of letters a occurring in a factor w , we may obtain the same information for $\varphi(w)$ by a simple formula. We mention the formula only for the binary alphabet $\mathcal{A} = \{0, 1\}$ we are interested in. It follows straightforwardly from the definition of the morphism matrix that for every factor $w \in \mathcal{A}^*$

$$(|\varphi(w)|_0, |\varphi(w)|_1) = (|w|_0, |w|_1)M_\varphi. \quad (2)$$

where $|v|_a$ denotes the number of letters a occurring in a word v . Clearly, a similar formula holds for any finite alphabet.

A morphism is said to be *primitive* if a power of M_φ has all elements strictly positive. In other words, matrices of primitive morphisms fulfill the assumptions of the Perron-Frobenius theorem.

The action of the morphism φ may be naturally extended to an infinite word $u = u_0u_1u_2\dots$ by the prescription

$$\varphi(u_0u_1u_2\dots) = \varphi(u_0)\varphi(u_1)\varphi(u_2)\dots$$

An infinite word u is called a *fixed point* of φ if $\varphi(u) = u$. It is known that any fixed point of a primitive morphism is uniformly recurrent and that any left eigenvector corresponding to the dominant eigenvalue of M_φ is proportional to the densities of letters in any fixed point of φ .

3 Infinite words associated with β -integers

Here, we provide the description of infinite words u_β associated with quadratic non-simple Parry numbers in terms of fixed points of primitive morphisms. We keep the notation from our precedent Letter [1], where the number theoretical background on β -integers and associated infinite words u_β is at disposal. Nevertheless, to make this Letter self-contained, we will recall all notions needed for understanding of our results. In this section we also deduce some important properties of u_β , in particular, a transformation generating bispecial factors of u_β . Bispecial factors turn out to be essential for our main aim – determination of the maximal integer powers of factors and determination of the index of u_β .

A non-simple quadratic Parry number β is the larger root of $x^2 - (p+1)x + p - q$, where $p > q \geq 1$. Its Rényi expansion of unity is $d_\beta(1) = pq^\omega$. The set of β integers has two distances between neighbors: $\Delta_0 = 1$ and $\Delta_1 = \beta - p$. Consequently, the infinite word u_β coding distances between neighboring β -integers is binary.

As we have already said, the reader may find the notions of the Rényi expansion of unity, β -integers, distances between neighbors in $\mathbb{Z}_\beta$ etc. in our precedent Letter [1]. However, in order to follow the ideas in the sequel, it is sufficient to know that u_β is the unique fixed point of a morphism canonically associated with parameters p, q characterizing non-simple quadratic Parry numbers β . Therefore we will use the result of [14] for an equivalent definition of u_β .

Definition 3.1. *Let β is the larger root of $x^2 - (p+1)x + p - q$, where $p > q \geq 1$. The unique fixed point of the morphism*

$$\varphi(0) = 0^p 1, \quad \varphi(1) = 0^q 1 \quad (3)$$

will be denoted u_β .

The infinite word u_β starts as follows

$$u_\beta = \underbrace{0^p 1 \dots 0^p 1}_{p \text{ times}} 0^q 1 \underbrace{0^p 1 \dots 0^p 1}_{p \text{ times}} 0^q 1 \dots \underbrace{0^p 1 \dots 0^p 1}_{p \text{ times}} 0^q 1 \underbrace{0^p 1 \dots 0^p 1}_{q \text{ times}} 0^q 1 \dots \quad (4)$$

The morphism matrix M_φ is $\begin{pmatrix} p & 1 \\ q & 1 \end{pmatrix}$ and φ is thus obviously primitive. Computing the left eigenvector of M_φ corresponding to the dominant eigenvalue β , we get the densities $1 - \frac{1}{\beta}$ and $\frac{1}{\beta}$ of letters 0 and 1, respectively.

Remark 3.2. *In the paper [10], it is shown that the factor complexity $\mathcal{C}$ of u_β satisfies*

$$\text{if } p > q + 1, \quad \text{then } \{\mathcal{C}(n+1) - \mathcal{C}(n) \mid n \in \mathbb{N}\} = \{1, 2\},$$

$$\text{if } p = q + 1, \quad \text{then } \{\mathcal{C}(n+1) - \mathcal{C}(n) \mid n \in \mathbb{N}\} = \{1\}.$$

Therefore, u_β is Sturmian if and only if $p = q + 1$.

First of all, some simple, but very important properties of the morphism φ are observed.

Observation 3.3. *Let $10^{k_1}1$ be a factor of u_β , then $k = p$ or $k = q$.*

Observation 3.4. *Let v be any factor of u_β containing at least one 1. Then there exists k_1 , $0 \leq k_1 \leq p$, such that $0^{k_1}1$ is a prefix of v and there exists k_2 , $0 \leq k_2 \leq p$, such that 10^{k_2} is a suffix of v . The fact that $\varphi(0)$ and $\varphi(1)$ end in 1 and contain only one letter 1 implies that there exists a unique word w in $\{0, 1\}^*$ satisfying $v = 0^{k_1}1\varphi(w)0^{k_2}$. Clearly, w is a factor of u_β .*

Of significant importance is the map $T : \{0, 1\}^* \rightarrow \{0, 1\}^*$ defined by

$$T(w) = 0^q 1 \varphi(w) 0^q. \quad (5)$$

The map T helps to generate bispecial factors that play a crucial role in the determination of the index of factors. Therefore, the rest of this section is devoted to the description of properties of T .

Lemma 3.5. *Let T be the map defined in (5).*

1. *For every $w \in \mathcal{L}(u_\beta)$, it holds that $T(w) \in \mathcal{L}(u_\beta)$.*
2. *Let w be a factor of u_β and let $a, b \in \mathcal{A}$, then $awb \in \mathcal{L}(u_\beta)$ if and only if $aT(w)b \in \mathcal{L}(u_\beta)$.*
3. *Let v be a bispecial factor of u_β containing at least one letter 1, then there exists a unique factor w such that $v = T(w)$.*
4. *Let w, v be factors of u_β , then w is a prefix of v if and only if $T(w)$ is a prefix of $T(v)$.*
5. *Let w, v be factors of u_β , then w is a suffix of v if and only if $T(w)$ is a suffix of $T(v)$.*

Proof. 1. Take an arbitrary factor $w \in \mathcal{L}(u_\beta)$. Then w is extendable to the right, and, since u_β is recurrent, w is also extendable to the left. In other words, there exists $a, b \in \{0, 1\}$ such that awb is also a factor of u_β . As u_β is a fixed point of φ , the image $\varphi(awb)$ belongs to $\mathcal{L}(u_\beta)$. Finally, $T(w)$ is a factor of u_β because $T(w)$ is a subword of $\varphi(awb)$.

2. Let $1w1$ be a factor of $\mathcal{L}(u_\beta)$, then $01w1$ is as well a factor of u_β . Applying φ , we learn that $\varphi(01w1) = \varphi(0)T(w)1$ is a factor of u_β , which proves that $1T(w)1$ belongs to $\mathcal{L}(u_\beta)$. The other cases $0w0, 0w1, 1w0$ are analogous.

Let $0T(w)1 \in \mathcal{L}(u_\beta)$, using Observation 3.3, the word $v = 10^p 1 \varphi(w) 0^q 1$ is also a factor of u_β . Applying Observation 3.4, $v = 1\varphi(0w1)$ and $0w1$ is an element of $\mathcal{L}(u_\beta)$. All the other cases $0T(w)0, 1T(w)0, 1T(w)1$ are similar.

3. Observation 3.3 implies that each bispecial factor v containing at least one letter 1 has the prefix $0^q 1$ and the suffix 10^q . According to Observation 3.4, there exists a unique w such that $v = T(w)$.

4. The implication $\Rightarrow$ is obvious noticing that 0^q is a prefix of $\varphi(a)$ for $a \in \{0, 1\}$. The opposite implication $\Leftarrow$ follows taking into account that $\varphi(1)$ is not a prefix of $\varphi(0)$ and $\varphi(0)$ is not a prefix of $\varphi(1a)$ for any $a \in \{0, 1\}$.

5. The implication $\Rightarrow$ is obvious noticing that $0^q 1$ is a suffix of $\varphi(a)$ for $a \in \{0, 1\}$. The opposite implication $\Leftarrow$ follows taking into account that $1\varphi(1)$ is not a suffix of $\varphi(0)$ and $\varphi(0)$ is not a suffix of $\varphi(x1)$ for any $x \in \{0, 1\}^*$. $\square$

4 Integer powers in u_β

Even if we want to describe the maximal integer powers of factors of u_β , it turns out to be useful to study first the relation between bispecial factors and the maximal rational powers of factors.

Lemma 4.1. *Let u be an infinite uniformly recurrent word over an alphabet $\mathcal{A}$. Let $w^k w'$ be its factor for some proper prefix w' of w and some positive integer k . Let us denote by **P1**, **P2**, **P3** the following statements:*

P1 *The factor w has the maximal index in u among all factors of u with the same length $|w|$ and $w^k w'$ is the maximal power of w in u .*

P2 *There exist $a, b \in \{0, 1\}$ such that*

$$aw^k w' b \in \mathcal{L}(u) \quad \text{and} \quad w' b \text{ is not a prefix of } w \quad \text{and} \quad a \text{ is not a suffix of } w.$$

P3 *All the following factors are bispecial:*

$$w', ww', www', \dots, w^{k-1} w'.$$

*Then **P1** implies **P2** and **P2** implies **P3**.*

Proof. **P1** $\Rightarrow$ **P2** : As u is recurrent, there exists a such that aw^kw' is a factor of u . Since w^kw' is the maximal power of w , the letter a is not a suffix of w , otherwise the factor awa^{-1} (usually called a conjugate of w) would have a larger index than w . On the other hand, if $w^kw'b$ is a factor of w , then $w'b$ is not a prefix of w , otherwise it contradicts the fact that w^kw' is the maximal power of w in u .

P2 $\Rightarrow$ **P3** : Since w' is a proper prefix of w , there exists $x \in \mathcal{A}$ such that $w'x$ is a prefix of w . Denote by y the last letter of w . Obviously, $x \neq b$ and $y \neq a$. As $aw^jw'x$ is a prefix of $aw^kw'b$ and $yw^jw'b$ is a suffix of $aw^kw'b$, both $aw^jw'x$ and $yw^jw'b$ are in $\mathcal{L}(u)$ for all j , $0 \leq j \leq k-1$. It follows that all factors listed in **P3** are bispecial. $\square$

In this section, our aim is to describe the maximal integer powers occurring in u_β . Since the letter 0 has the maximal index p , we may restrict our consideration to k -th powers of factors of u_β with $k \geq p$. Crucial for the determination of the index of u_β are Propositions 4.2 and 4.5.

Proposition 4.2. *Let p, q be integers, $p > q \geq 1$, and u_β be the fixed point of the morphism (3). Assume $p > 3$. Let w be a factor of u_β containing at least two 1s and w' be a proper prefix of w . Denote $v = w^kw'$ for some $k \in \mathbb{N}$, $k \geq p$. If there exist $a, b \in \{0, 1\}$ so that*

$$avb \in \mathcal{L}(u_\beta) \quad \text{and} \quad w'b \text{ is not a prefix of } w \quad \text{and} \quad a \text{ is not a suffix of } w,$$

then there exist a unique $\tilde{w}$ of length ≥ 2 and a proper prefix $\tilde{w}'$ of $\tilde{w}$ such that

$$w = 0^q 1 \varphi(\tilde{w})(0^q 1)^{-1} \quad \text{and} \quad v = T(\tilde{v}) = T(\tilde{w}^k \tilde{w}'); \quad (6)$$

moreover,

$$a\tilde{w}b \in \mathcal{L}(u_\beta) \quad \text{and} \quad \tilde{w}'b \text{ is not a prefix of } \tilde{w} \quad \text{and} \quad a \text{ is not a suffix of } \tilde{w}.$$

In order to prove Proposition 4.2, we will use the following lemma.

Lemma 4.3. *Let p, q be integers, $p > q \geq 1$, and u_β be the fixed point of the morphism (3). The following statements hold:*

1. *If $0(x1)^\ell x0 \in \mathcal{L}(u_\beta)$ for some integer $\ell \geq 2$, then $\ell = 2$ and $x = 0^q$.*
2. *If $1(x0)^\ell x1 \in \mathcal{L}(u_\beta)$ for some integer $\ell \geq p-1$ and $p \leq 2q$, then x is the empty word ε .*

Proof. 1. At first, we exclude the case when x contains a non-zero letter. Suppose that the letter 1 occurs in x . Since the factors $0x1$ and $1x0$ belong to $\mathcal{L}(u_\beta)$, it follows that x is bispecial. By Lemma 3.5 Item 3., x starts in $0^q 1$ and ends in 10^q . Therefore $10^q 10^q 1 \in \mathcal{L}(u_\beta)$. As $10^q 10^q 1 = 1\varphi(11)$, we have according to Observation 3.4 that $11 \in \mathcal{L}(u_\beta)$ – a contradiction.

Now consider $x = 0^s$ for some $s \in \mathbb{N}$. Then $0x1, 1x1 \in \mathcal{L}(u_\beta)$, which implies by Observation 3.3 that $s = q$. If ℓ was at least 3, then $1x1x1 = 10^q 10^q 1 \in \mathcal{L}(u_\beta)$, which leads to the same contradiction as before.

2. Again we start with the case of x containing the letter 1. Since factors $1x0$ and $0x1$ belong to the language $\mathcal{L}(u_\beta)$, it follows that x is bispecial. Hence x starts in $0^q 1$ and ends in 10^q . Since $10^q 00^q 1 \in \mathcal{L}(u_\beta)$, by Observation 3.3, we have $p = 2q + 1$. This contradicts the assumption $p \leq 2q$.

Suppose now that $x = 0^s$ for some $s \in \mathbb{N}$. Since $1(x0)^\ell x1 = 1(0^{s+1})^\ell 0^s 1$ is a factor of u_β , Observation 3.3 gives that $(s+1)\ell + s \leq p$, which is impossible if $s \geq 1$ and $\ell \geq p-1$. Therefore $s = 0$ and x is the empty word. $\square$

Proof of Proposition 4.2. The factor w contains at least two 1s. Since w and $v = w^kw'$ satisfy Item **P2** of Lemma 4.1, both ww' and www' are bispecial, and therefore start in $0^q 1$ and end in 10^q . Consequently, their form is $ww' = T(x)$ and $www' = T(y)$, where $x, y \neq \varepsilon$. According to Lemma 3.5 Item 5., x is a suffix of y , i.e., $y = zx$ for some $z \neq \varepsilon$. Observing $ww' = 0^q 1 \varphi(x) 0^q$ and $www' = 0^q 1 \varphi(z) \varphi(x) 0^q$, it follows directly that $w = 0^q 1 \varphi(z) (0^q 1)^{-1}$.

Let us, at first, show that

1. either z is a prefix of x ,
2. or $z = x1$,
3. or $z = x0$.

Assume $z = tdz'$ and $x = t(1-d)x'$ for a word $t \in \{0,1\}^*$ and for a letter d . Then $w = 0^q 1 \varphi(t) \varphi(d) \varphi(z') (0^q 1)^{-1}$ is a prefix of $ww' = 0^q 1 \varphi(t) \varphi(1-d) \varphi(x') 0^q$. If $z' \neq \varepsilon$, we have a contradiction immediately. If $z' = \varepsilon$, then $t \neq \varepsilon$ knowing that z contains at least two letters (w contains at least two 1s).

- If $d = 1$, then $w = 0^q 1 \varphi(t)$ and $ww' = 0^q 1 \varphi(t) 0^p 1 \varphi(x') 0^q$, thus w' starts in $0^p 1$, which is not a prefix of w – a contradiction.
- If $d = 0$, then $w = 0^q 1 \varphi(t) 0^{p-q}$ and $ww' = 0^q 1 \varphi(t) 0^{p-q} 0^{2q-p} 1 \varphi(x') 0^q$, thus w' starts in $0^{2q-p} 1$, which is not a prefix of w because $p \neq q$ – a contradiction again.

The situation $z = xz'$ for some z' of length ≥ 2 cannot occur because it implies $|w| > |ww'|$. Consequently, one of the situations 1., 2., 3. occurs.

1. If z is a prefix of x , i.e., $x = zx''$, then $ww' = 0^q 1 \varphi(z) \varphi(x'') 0^q$ and $w = 0^q 1 \varphi(z) (0^q 1)^{-1}$, thus $w' = 0^q 1 \varphi(x'') 0^q$. Then $v = w^k w' = 0^q 1 \varphi(z^k x'') 0^q$, therefore $\tilde{w} = z$, $\tilde{w}' = x''$. As w' is a proper prefix of w , it follows by Lemma 3.5 Item 4 that $\tilde{w}'$ is a proper prefix of $\tilde{w}$.
2. Assume $z = x1$, then $w = 0^q 1 \varphi(x)$ and $ww' = 0^q 1 \varphi(x) 0^q$, thus $w' = 0^q$. Then $v = 0^q 1 \varphi((x1)^{k-1} x) 0^q$. Since $w'0$ is not a prefix of w and 0 is not a suffix of w , the assumptions imply that $0v0 \in \mathcal{L}(u_\beta)$. By Observation 3.3, we have $1\varphi(0(x1)^{k-1} x0) \in \mathcal{L}(u_\beta)$. Since $k-1 \geq 3$, we deduce by Lemma 4.3 that $0(x1)^{k-1} x0$ is not a factor of u_β . It contradicts Observation 3.4. Hence, the case $z = x1$ does not occur.
3. If $z = x0$, then $w = 0^q 1 \varphi(x) 0^{p-q}$ and $ww' = 0^q 1 \varphi(x) 0^q$, thus $w' = 0^{2q-p}$, which can happen only for $p \leq 2q$. Then $v = 0^q 1 \varphi((x0)^{k-1} x) 0^q$. Since $w'1$ is not a prefix of w and 1 is not a suffix of w , the assumptions imply that $1v1 \in \mathcal{L}(u_\beta)$. Hence, $1\varphi(1(x0)^{k-1} x1) \in \mathcal{L}(u_\beta)$. However, by Lemma 4.3, it follows that $x = \varepsilon$. Then $z = 0$, which contradicts the condition $|z| \geq 2$. Thus, the case $z = x0$ does not occur.

Let us finally note that the very last statement on the extensions of $\tilde{v}$ follows from Lemma 3.5 Items 2., 4., and 5. $\square$

Remark 4.4. Proposition 4.2 does not take into account u_β given by parameters $p = 2, q = 1$, $p = 3, q = 2$, and $p = 3, q = 1$. In the two first cases, u_β is a Sturmian word. Therefore, exclusion of the first two cases does not mean any loss. For the case of $p = 3, q = 1$, in the proof of Proposition 4.2, we cannot exclude the situation 2.; in this case, Lemma 4.3 Item 1. implies either the validity of (6) or of

$$w = 01\varphi(01)(01)^{-1} \quad v = T(01010).$$

Proposition 4.2 thus claims that for every factor $w \in \mathcal{L}(u_\beta)$ containing at least two 1s such that its k -th power w^k is a factor of u_β with $k \geq p$, there exists a shorter factor $\tilde{w}$ such that its k -th power $\tilde{w}$ is also in the language of u_β . As a consequence, in order to determine the maximal integer power present in u_β , it is sufficient to study the index of factors w containing only one letter 1.

Proposition 4.5. Let p, q be integers, $p > q \geq 1$, and u_β be the fixed point of the morphism (3). Let w be a factor of u_β containing one letter 1 and of the maximal index $\text{ind}(w)$ among all factors of length $|w|$ and such that $\text{ind}(w) \geq p \geq 3$. Denote $k := \lfloor \text{ind}(w) \rfloor$ and $v = w^k w'$ the maximal power of w . Then

$$w = 0^q 1 \varphi(0)(0^q 1)^{-1} \quad \text{and} \quad v = T(0^p)$$

and

$$\text{ind}(w) = p + \frac{2q+1}{p+1}.$$

Proof. According to Lemma 4.1, ww' is a bispecial factor. Lemma 3.5 Item 3. claims that ww' starts in $0^q 1$. Therefore $w = 0^q 10^s$ with $s \in \{0, p - q\}$ (Observation 3.3). The case $s = 0$ does not occur since $www = 0^q 10^s 0^q 10^s 0^q 10^s \in \mathcal{L}(u_\beta)$ and $0^q 10^q 10^q 1 = 0^q 1 \varphi(11)$, but $11 \notin \mathcal{L}(u_\beta)$ – a contradiction to Observation 3.4. Hence $w = 0^q 10^{p-q} = 0^q 1 \varphi(0)(0^q 1)^{-1}$. It remains to determine the form of v . Again, since ww' is bispecial, ww' ends in 10^q . As w' is a prefix of w , at most one 1 occurs in w' .

- Suppose w' contains 1, then w' starts in $0^q 1$ and ends in 10^q , thus $w' = 0^q 10^q$. This is possible only in case when $p - q - 1 \geq q$, i.e., $p \geq 2q + 1$. Then, $v = (0^q 10^{p-q})^k 0^q 10^q$. Consequently, $v = 0^q 1 \varphi(0^k) 0^q = T(0^k)$. On one hand, Observations 3.3 and 3.4 imply that $k \leq p$. On the other hand, since v is the maximal power of w , it follows that $k \geq p$.
- Assume w' does not contain 1. Since ww' ends in 10^q and $w = 0^q 10^{p-q}$, the only possibility for w' is $w' = 0^{2q-p}$. This comes in question only for $p \leq 2q$. Then $v = (0^q 10^{p-q})^k 0^{2q-p} = 0^q 1 \varphi(0^{k-1}) 0^q = T(0^{k-1})$. The same arguments as in the previous case imply that $k - 1 = p$.

Clearly, $\text{ind}(w) = \frac{|v|}{|w|} = p + \frac{2q+1}{p+1}$. $\square$

Let us state the main result of this section.

Theorem 4.6. *Let p, q be integers, $p > q \geq 1$, and u_β be the fixed point of the morphism (3). Assume $p \geq 3$.*

- *If $p \leq 2q$, then there exists a factor $w \neq \varepsilon$ satisfying $w^{p+1} \in \mathcal{L}(u_\beta)$ and no $(p+2)$ -nd power of any factor belongs to the language $\mathcal{L}(u_\beta)$.*
- *If $p > 2q$, then there exists a factor $w \neq \varepsilon$ satisfying $w^p \in \mathcal{L}(u_\beta)$ and no $(p+1)$ -st power of any factor belongs to the language $\mathcal{L}(u_\beta)$.*

Proof. Proposition 4.2 implies that in order to determine the maximal integer power present in u_β , we can restrict our consideration to powers of factors containing only one letter 1. When we compute the integer part $\lfloor \text{ind}(w) \rfloor$ of such factors w in Proposition 4.5, we find that the maximum is $p+1$ if $p \leq 2q$ and p otherwise. $\square$

5 Index of u_β

The task of this section is to compute the index of u_β , i.e.,

$$\text{ind}(u_\beta) = \sup\{\text{ind}(w) \mid w \in \mathcal{L}(u_\beta)\}.$$

We already know that $\text{ind}(u_\beta) \geq p$. Using Lemma 4.1, it suffices to study rational powers $v = w^k w'$ of factors w with the property **P2**. As a direct consequence of Propositions 4.2 and 4.5, we have the following corollary.

Corollary 5.1. *Let p, q be integers, $p > q \geq 1$, and u_β be the fixed point of the morphism (3). Assume $p > 3$. The index of u_β is given by the following formula*

$$\text{ind}(u_\beta) = \sup\{\text{ind}(w^{(n)}) \mid n \in \mathbb{N}\},$$

where

$$w^{(0)} = 0, \quad w^{(n+1)} = 0^q 1 \varphi(w^{(n)}) (0^q 1)^{-1}. \quad (7)$$

Moreover, the maximal power of $w^{(n)}$ is $v^{(n)}$, where

$$v^{(0)} = 0^p, \quad v^{(n+1)} = T(v^{(n)}). \quad (8)$$

In the sequel, let us determine the index of $w^{(n)}$ for every $n \in \mathbb{N}$.

Lemma 5.2. *The number of 0s and 1s in the words $w^{(n)}$ and $v^{(n)}$ satisfy*

$$\begin{aligned} (|w^{(n)}|_0, |w^{(n)}|_1) &= (1, 0)M_\varphi^n, \\ (|v^{(n)}|_0, |v^{(n)}|_1) &= (p+1, \frac{2q+1-p}{q})M_\varphi^n - (1, \frac{2q+1-p}{q}), \end{aligned}$$

where $M_\varphi = \begin{pmatrix} p & 1 \\ q & 1 \end{pmatrix}$ is the morphism matrix.

Proof. As $w^{(n)}$ is a conjugate of $\varphi(w^{(n-1)})$, the first formula holds by (2). Let us show the second one by induction on n .

For $n = 0$,

$$(|v^{(0)}|_0, |v^{(0)}|_1) = (p, 0) = (p+1, \frac{2q+1-p}{q}) - (1, \frac{2q+1-p}{q}).$$

For $n > 0$,

$$(|v^{(n)}|_0, |v^{(n)}|_1) = (|v^{(n-1)}|_0, |v^{(n-1)}|_1)M_\varphi + (2q, 1) =$$

by the induction assumption,

$$= \left[(p+1, \frac{2q+1-p}{q})M_\varphi^{n-1} - (1, \frac{2q+1-p}{q}) \right] M_\varphi + (2q, 1) = (p+1, \frac{2q+1-p}{q})M_\varphi^n - (1, \frac{2q+1-p}{q}).$$

□

Since the eigenvalues β and β' of M_φ are roots of the Parry polynomial $x^2 - (p+1)x + (p-q)$, it is straightforward to show that $\vec{x}_1 = (\beta - 1, 1)$ is a left eigenvector of M_φ corresponding to β and $\vec{x}_2 = (\beta' - 1, 1)$ is a left eigenvector of M_φ corresponding to β' . The index of $w^{(n)}$ may be expressed as follows

$$\begin{aligned} \text{ind}(w^{(n)}) &= \frac{|v^{(n)}|}{|w^{(n)}|} = \frac{(p+1, 0)M_\varphi^n \begin{pmatrix} 1 \\ 1 \end{pmatrix} + (0, \frac{2q+1-p}{q})M_\varphi^n \begin{pmatrix} 1 \\ 1 \end{pmatrix} - (1, \frac{2q+1-p}{q}) \begin{pmatrix} 1 \\ 1 \end{pmatrix}}{(1, 0)M_\varphi^n \begin{pmatrix} 1 \\ 1 \end{pmatrix}}, \\ \text{ind}(w^{(n)}) &= p+1 + \frac{\frac{2q+1-p}{q}(\alpha_1 \vec{x}_1 + \alpha_2 \vec{x}_2)M_\varphi^n \begin{pmatrix} 1 \\ 1 \end{pmatrix} - \frac{3q+1-p}{q}}{(\gamma_1 \vec{x}_1 + \gamma_2 \vec{x}_2)M_\varphi^n \begin{pmatrix} 1 \\ 1 \end{pmatrix}}, \end{aligned}$$

where $\alpha_1 \vec{x}_1 + \alpha_2 \vec{x}_2 = (0, 1)$ and $\gamma_1 \vec{x}_1 + \gamma_2 \vec{x}_2 = (1, 0)$. Using the fact that $\vec{x}_1$ and $\vec{x}_2$ are eigenvectors of M_φ , we have

$$\text{ind}(w^{(n)}) = p+1 + \frac{\frac{2q+1-p}{q}(\alpha_1 \beta^n \vec{x}_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \alpha_2 \beta'^n \vec{x}_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}) - \frac{3q+1-p}{q}}{\gamma_1 \beta^n \vec{x}_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \gamma_2 \beta'^n \vec{x}_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}}.$$

It is easy to calculate that $\alpha_1 = \frac{1-\beta'}{\beta-\beta'}$, $\alpha_2 = \frac{\beta-1}{\beta-\beta'}$, $\gamma_1 = \frac{1}{\beta-\beta'}$, $\gamma_2 = \frac{-1}{\beta-\beta'}$, and $\vec{x}_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \beta$, $\vec{x}_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \beta'$. The final formula for the index of $w^{(n)}$ has the form

$$\begin{aligned} \text{ind}(w^{(n)}) &= p+1 + \frac{\frac{2q+1-p}{q}((1-\beta')\beta^{n+1} - (1-\beta)\beta'^{n+1}) - \frac{3q+1-p}{q}(\beta - \beta')}{\beta^{n+1} - \beta'^{n+1}} = \\ &= p+1 + \frac{2q+1-p}{q}(1-\beta') + \underbrace{\frac{\beta - \beta'}{q(\beta^{n+1} - \beta'^{n+1})}}_{>0} \underbrace{\left((2q+1-p)\beta'^{n+1} - (3q+1-p) \right)}_{A(n)}. \end{aligned}$$

Using the fact $0 < \beta' < 1 < \beta$, we determine the limit

$$\lim_{n \rightarrow \infty} \text{ind}(w^{(n)}) = p+1 + \frac{2q+1-p}{q}(1-\beta') = p+1 + \frac{2q+1-p}{\beta-1}.$$

This limit is the supreme of $\{\text{ind}(w^{(n)}) \mid n \geq 0\}$ if and only if $A(n) < 0$ for all $n \in \mathbb{N}$. It is an easy exercise to show that $A(n) = (2q+1-p)(\beta'^{n+1} - 1) - q < 0$ for all $n \in \mathbb{N}$ if and only if $p \leq 3q+1$, otherwise $A(n) > 0$ for all sufficiently large n .

Let us sum up the results in a theorem.

Theorem 5.3. *Let p, q be integers, $p > q \geq 1$, and u_β be the fixed point of the morphism (3). Assume $p > 3$. Then the index of u_β satisfies for $p \leq 3q + 1$*

$$\text{ind}(u_\beta) = p + 1 + \frac{2q + 1 - p}{\beta - 1},$$

otherwise there exists $n_0 \in \mathbb{N}$ such that

$$\text{ind}(u_\beta) = \text{ind}(w^{(n_0)}) > p + 1 + \frac{2q + 1 - p}{\beta - 1}.$$

Remark 5.4. *Similarly as in the previous section, we have to treat the case of $p = 3$ and $q = 1$ separately. According to Remark 4.4, we have to determine the index of $(w^{(n)})$ defined in (7), but moreover the index of $(\hat{w}^{(n)})$ defined recursively by*

$$\hat{w}^{(0)} = 01\varphi(01)(01)^{-1}, \quad \hat{w}^{(n+1)} = 0^q 1 \varphi(\hat{w}^{(n)})(0^q 1)^{-1}.$$

Using the same technique as before, we obtain

$$\sup\{\text{ind}(\hat{w}^{(n)}) \mid n \in \mathbb{N}\} = \beta < 4 = \sup\{\text{ind}(w^{(n)}) \mid n \in \mathbb{N}\}.$$

Hence, Theorem 5.3 holds in fact also in this case.

At the conclusion, let us compare in case of Sturmian words u_β the formula for $\text{ind}(w^{(n)})$ with the formula (1) for the index of general Sturmian words. As we have already stated, u_β is Sturmian if and only if $p = q + 1$, i.e., β is the larger root of the polynomial $x^2 - (p + 1)x + 1$. For such β , we have

$$\text{ind}(u_\beta) = p + 1 + \frac{2p - 1}{\beta - 1} = \beta + 1.$$

In order to apply the formula from (1), we need to determine the slope α of the Sturmian word u_β . Since $\alpha > \frac{1}{2}$ is the density of the more frequent letter, according to Section 3, $\alpha = 1 - \frac{1}{\beta}$. Let us use some basic properties of continued fractions available at any book on Number Theory to determine the continued fraction of this value. Since

$$1 - \frac{1}{\beta} = \frac{1}{1 + \frac{1}{\beta - 1}} = \frac{1}{1 + \frac{1}{p - 1 + 1 - \frac{1}{\beta}}},$$

one obtains $\beta = [0; 1, (p - 1), 1, (p - 1), \dots] = [0; \overline{1, (p - 1)}]$. Denominators q_n of the convergents of β fulfill therefore the following recurrent relations

$$q_{2n+1} = (p - 1)q_{2n} + q_{2n-1} \quad \text{and} \quad q_{2n} = q_{2n-1} + q_{2n-2}$$

with initial values $q_1 = 1$, $q_2 = p$, $q_3 = p + 1$. By mathematical induction on n , it may be shown easily that

$$q_{2n-1} = \frac{1}{\beta - \beta'} (\beta^n - \beta'^n) \quad \text{and} \quad q_{2n} = \frac{1}{\beta - \beta'} ((1 - \beta')\beta^{n+1} - (1 - \beta)\beta'^{n+1}).$$

As it holds for coefficients of the continued fraction of β that $a_{2n-1} = 1$ and $a_{2n} = p - 1$, it suffices to consider even n in (1). We obtain then finally

$$a_{2n+2} + 2 + \frac{q_{2n} - 2}{q_{2n+1}} = p + 1 + \frac{((1 - \beta')\beta^{n+1} - (1 - \beta)\beta'^{n+1}) - 2(\beta - \beta')}{\beta^{n+1} - \beta'^{n+1}},$$

which is exactly $\text{ind}(w^{(n)})$. This result holds for all parameters p, q satisfying $p = q + 1$, even for $p \leq 3$. Consequently, Theorem 5.3 is in fact valid for all parameters p, q with $p > q \geq 1$.

Acknowledgements

The authors acknowledge financial support by Czech Science Foundation GA ĆR 201/05/0169, by the grants MSM6840770039 and LC06002 of the Ministry of Education, Youth, and Sports of the Czech Republic.

References

- [1] Balková, Ĺ., Gazeau, J.-P., Pelantová, E.: Asymptotic behavior of beta-integers, *Letters in Mathematical Physics* 84 179-198 (2008)
- [2] Frougny, Ch., Gazeau, J.-P., Krejcar, R.: Additive and multiplicative properties of point-sets based on beta-integers, *Theor. Comp. Sci.* 303 491-516 (2003)
- [3] Berstel, J.: On the index of Sturmian words, in: *Jewels are forever*, Springer 287-294 (1999)
- [4] Justin, J., Pirillo, G.: Fractional powers in Sturmian words, *Theoret. Comput. Sci.* 255 363–376 (2001)
- [5] Mignosi, F., Pirillo, G.: Repetitions in the Fibonacci infinite word, *RAIRO Inform. Theor. Appl.* 26 199–204 (1992)
- [6] Masáková, Z., Pelantová, E.: Relation between powers of factors and recurrence function characterizing Sturmian words, submitted, arXiv 0809.0603v2[math.CO]
- [7] Carpi, A., de Luca, A.: Special factors, periodicity, and an application to Sturmian words, *Acta Inform.* 36 983-1006 (2000)
- [8] Damanik, D., Lenz, D.: The index of Sturmian sequences, *European J. Combin.* 23 23-29 (2002)
- [9] Damanik, D.: Singular continuous spectrum for a class of substitution Hamiltonians II, *Letters in Mathematical Physics* 54 25-31 (2000)
- [10] Balková, Ĺ., Masáková, Z.: Palindromic complexity of infinite words associated with non-simple Parry numbers, *RAIRO - Theoret. Informatics Appl.* 43 145-163 (2009)
- [11] Balková, Ĺ., Pelantová, E., Turek, O.: Combinatorial and arithmetical properties of infinite words associated with quadratic non-simple Parry numbers, *RAIRO - Theoret. Informatics Appl.* 41 307-328 (2007)
- [12] Berstel, J.: Sturmian and episturmian words (a survey of some recent results), in: S. Bozapalidis, G. Rahonis (eds), *Conference on Algebraic Informatics*, Thessaloniki, *Lecture Notes Comput. Sci.* 4728 23-47 (2007)
- [13] Lothaire, M.: *Algebraic combinatorics on words*, volume 90 of *Encyclopedia of Mathematics and its Applications*, Cambridge University Press (2002)
- [14] Fabre, S.: Substitutions et β -systèmes de numération, *Theoret. Comput. Sci.* 137 219-236 (1995)